


# 2021 STI Treatment Guidelines

Hillary Liss, MD  
Clinical Associate Professor, University of WA  
August 25, 2021



# Disclosure

---

- Hillary Liss has no relevant financial relationships with an entity producing, marketing, re-selling, or distributing health care goods or services consumed by, or used on patients.

Syphilis management? Resistant gonorrhea? STD treatment?

# GOT A TOUGH STD QUESTION?

Get FREE expert STD clinical consultation at your fingertips




Ask your question

National STD  
experts review

Response within 1-5  
business days,  
depending on urgency

Log on to **[www.STDCCN.org](http://www.STDCCN.org)** for medical professionals nationwide


# 2019 was worst year on record for reported STIs




OVER HALF OCCURRED AMONG YOUNG PEOPLE 15-24 years of age



# New guidelines



- Hot off the presses!
- July 23, 2021



- Expected later in 2021

# What's in a name?

## — STD

- Sexually transmitted disease
- Refers to disease state




## vs STI

- Sexually transmitted infection
- Refers to pathogen
- Often asymptomatic



# Evidence-based Approach to Guideline Development



# CDC STI Treatment Guideline Development

---

- Evidence-based on principal outcomes of STI therapy
- “Recommended” regimens preferred over “alternative” regimens
- Treatments alphabetized unless there is a priority of choice
- Released July 2021
  - Available at: <https://www.cdc.gov/std/treatment-guidelines/toc.htm>
  - Interim app download: <https://www.cdc.gov/STIapp/>





# Screening



# STI Screening for Cis-Women (WSM and WSW)

## Women under 25 years of age

Chlamydia/gonorrhea

HIV at least once

Hep C at least once if  $\geq 18$  yo (unless prevalence of Hep C  $< 0.1\%$ )

## Women 25 years of age and older

Chlamydia/gonorrhea if at risk

HIV at least once

Hep C at least once (unless prevalence of Hep C  $< 0.1\%$ )

## Pregnant women

Chlamydia (<25 years of age or risk and retest during 3<sup>rd</sup> trimester)

Gonorrhea (<25 years of age or risk and retest during 3<sup>rd</sup> trimester)

HIV

Syphilis serology

HepB sAg

Hep C (unless prevalence of Hep C  $< 0.1\%$ ) WITH EVERY PREGNANCY



# STI Screening for Transgender Persons

## Based on current anatomy and gender of sex partners

- Offer HIV screening to all transgender persons
- TG persons who have sex with cisgender men, at similar risk for STIs as cis-MSM

## Transgender women post vaginoplasty

- GC/CT (all sites of exposure: oral, anal, genital)

(Urine vs neovaginal swab not specified, best specimen type based on tissue type used to construct neovagina)

## Transgender Men post metoidioplasty

- If vagina still present and need to screen for STIs, cervical (or vaginal) swab should be used



# STI Screening for cis-MSM

- HIV\*
- Syphilis\*
- Urethral GC and CT\*
- Rectal GC and CT (if receptive anal sex)\*
- Pharyngeal GC (if oral sex)\*
- Hepatitis B (HBsAg, HBV core ab, HBV surface ab)
- Hepatitis C: (At least once if  $\geq 18$  yo, unless prevalence of infection  $< 0.1\%$ )
- Anal cancer: annual digital anorectal exam may be useful (no anal Pap rec yet)
- HSV-2 serology (consider)

- At least annually, more frequent (3-6 months) if multiple/anonymous partners, drug use, or partners w/ risk
- Routine screening not recommended for M. genitalium





# What about “Extragenital” Screening?

---

- Extragenital screening = testing for STIs at any body site other than genitourinary (urethral/urine/vaginal/cervix)
- Usually refers to rectal and oropharynx
- Typically for gonorrhea and/or chlamydia only
- Recommended routinely only for men who have sex with men (MSM), but now permissive for other individuals

# Importance of Extragenital GC/CT Infections

- Transmission
  - 30% of symptomatic gonococcal urethritis is attributable to oro-pharyngeal exposure<sup>1</sup>
- HIV Transmission
  - Can potentiate acquisition, even after controlling for sexual behaviors<sup>2-4</sup>
- Treatment can differ
  - Pharyngeal GC<sup>5</sup>
    - Ceftriaxone > Cefixime
  - Rectal CT<sup>6</sup>
    - Doxy >>> Azithromycin

# Extragenital Gonorrhea & Chlamydia is Common

- Among MSM, high rates of extra-genital GC & CT
  - Pharyngeal GC: 9.2%<sup>1</sup>
  - Rectal GC: 9.7%<sup>3</sup>
  - Rectal CT: 12%<sup>3</sup>
- The majority of infections are asymptomatic
  - 92% of pharyngeal GC<sup>2</sup>
  - 84 - 86% of rectal GC<sup>2</sup>

# Don't forget the triple dip: STD Screening for MSM



← Syphilis & HIV serology

← Pharyngeal GC


← Urine GC/CT

← Rectal GC/CT

Annually for all sexually active MSM  
Every 3-6 months for high-risk MSM



# STI Self-Testing Program



Available in English and Spanish



California PTC

Email [aradford@uw.edu](mailto:aradford@uw.edu) or go to <https://www.uwptc.org/visual-guides> for free posters for your clinic

# STI Screening for cis-MSW

---

- Routine STI screening not recommended
  - GC/CT recommended in high prevalence settings (e.g. adolescent clinics, correctional facilities, and STI clinics)
- HIV at least once (between 13-64 years of age)
- Hep C at least once if  $\geq 18$  yo (unless prevalence of Hep C  $< 0.1\%$ )

# Who should be Screened for CT/GC?

## Females

- < 25 annually, 25+ if at risk
- Pregnant <25 or risk

## MSM

- 3-6 month intervals at all exposed sites: genital, rectal, pharyngeal

## MSW

- High prevalence settings (e.g., Corrections, STI Clinics, adolescents)

## Persons living with HIV

- At least annually
- All exposed sites: genital, rectal, pharyngeal

## Patients on PrEP

- Every 3-6 months
- All exposed sites

## Adolescents

- Consider rectal/pharyngeal screen based on reported behavior/ exposure

# Who should be screened for Syphilis

## Pregnancy

- At first prenatal visit
- Again **at 28 weeks and at delivery** (if at high risk, or residing in area with high syphilis morbidity)

## MSM

- Including those on PrEP, 3-6 month intervals

## Corrections

- Universal **opt out** screening on intake based on local area or institutional incidence

## Persons living with HIV

- At least annually

## STI Clinic patients

- Regardless of symptoms
- If other STI diagnosed



# Who should be screened for HIV?

---

- CDC recommends: At least one time screening for all patients aged 13-64 years
  - All persons who seek STI screening
- USPSTF recommends:
  - Screen people aged 15 to 65 years
  - Risk-based screening for younger adolescents & older adults
  - Pregnant women regardless of age




# Chlamydia




# Doxycycline vs Azithromycin for Urogenital Chlamydia

## Efficacy




Slide credit: Dr. Will Geisler

# Doxycycline vs Azithromycin for Rectal Chlamydia



Slide credit: Dr. Will Geisler

# Randomized Controlled Trial DOX vs AZM for Rectal CT: Microbiologic Cure at 4 Weeks



Slide credit, J. Dombrowski

Dombrowski J, 2021, *CID* <https://doi.org/10.1093/cid/ciab153>

# Chlamydia Treatment:

## Urogenital/ Rectal/ Pharyngeal

Change in 2021 STI Treatment Guidelines

### Recommended regimens (non-pregnant):

- Doxycycline 100 mg orally twice daily for 7 days\*

### Alternative regimens (non-pregnant):

- Azithromycin 1 g orally in a single dose OR
- Levofloxacin 500 mg orally once daily for 7 days

\*Doxycycline delayed-release 200 mg, once-daily dosing for 7 days effective for urogenital CT. More costly but lower frequency GI side effects than standard doxycycline.

# Chlamydia Treatment: Pregnancy

## Recommended regimen (pregnant\*):

- Azithromycin 1 g orally in a single dose

## Alternative regimens (pregnant\*):

- Amoxicillin 500 mg orally three times a day for 7 days

**\* Test of cure at 3-4 weeks only in pregnancy**

# Expedited Partner Therapy for GC/CT

---

- No states in US prohibit EPT (either allowable or potentially allowable by law/statute in all 50 states)
- Previously only recommended for hetero men/women, now “shared decision making” for EPT for MSM
- Providing patients with packaged oral medications is preferred approach
  - Partners (especially adolescents) may not fill prescriptions



Sure feels like  
there are a lot of  
changes for me in  
the 2021 CDC STI  
Guidelines!



Chlamydia

Hold my  
beer...



Gonorrhea





# Gonorrhea



# More than half of GC isolates are resistant to at least one antibiotic

## Prevalence of Resistant or Decreased Susceptibility of *N. gonorrhoeae* Isolates to Antimicrobials, GISP, 2009 and 2019\*



\* 2019 data are preliminary

# **\*New\*** Gonorrhea Treatment Guidelines

for uncomplicated infections

Ceftriaxone **500** mg IM x 1  
for persons weighing <150kg\*

\*For persons weighing  $\geq 150$  kg, 1 g of IM  
ceftriaxone should be administered

However, if chlamydia has not been  
excluded, treat for chlamydia with:

Doxycycline 100 mg PO  
BID x 7 days

For pregnancy, allergy, or concern  
for non-adherence, 1 g PO  
azithromycin x 1 can be used

- No longer recommending dual therapy with azithromycin
- Test-of-Cure at 7-14 days post treatment for **pharyngeal** gonorrhea

# **\*New\*** **Alternative** Gonorrhea Treatment

for uncomplicated infections of the cervix, urethra, and rectum if ceftriaxone is not available:

Cefixime 800 mg PO x 1

However, if chlamydia has not been excluded, treat for chlamydia with:

Doxycycline 100 mg PO  
BID x 7 days

For pregnancy, allergy, or concern for non-adherence, 1 g PO azithromycin x 1 can be used

**Cephalosporin allergy: Gentamicin 240 mg IM + azithromycin 2 g PO**

No reliable alternative treatments are available for pharyngeal gonorrhea

# Rationale for GC Treatment Changes

---

- Improved antimicrobial stewardship
- Pharmacokinetic and pharmacodynamic considerations
- Changes in azithromycin susceptibility in GC

# Antimicrobial Stewardship

Need to minimize antibiotic exposure unless benefit outweighs risk

Risk benefit of dual vs monotherapy for GC

Drug-Resistant GC Urgent Threat

- Azithromycin resistance is a concern for other bacteria, so want to reduce overall use of azithromycin




# Pharmacodynamics/ Pharmacokinetics

---

- Antibiotic most effective when drug levels are above MIC ( minimum inhibitory concentration): lowest concentration of antibiotic needed to kill the bacteria
- Ceftriaxone kills GC when levels are high enough for long enough
  - 20-24 hours for Ceftriaxone
  - 500 mg dose most effective
  - Via modeling/mouse model
- Higher dose also more likely to kill gonorrhea in the pharynx




# Rise in GC Isolates with Decreased Susceptibility to Azithromycin (~5%) Gonococcal Isolate Surveillance Project (GISP), 2010–2019



**NOTE:** Elevated MIC = Azithromycin:  $\geq 2.0 \mu\text{g/mL}$ ; Cefixime:  $\geq 0.25 \mu\text{g/mL}$ ; Ceftriaxone:  $\geq 0.125 \mu\text{g/mL}$



# **Mycoplasma genitalium**



# More than 1 in 4 men with urethritis have *Mycoplasma genitalium*

## MAGNUM STUDY

Men with urethritis symptoms were enrolled from 6 U.S. STD clinics during 6/2017–8/2018


Study Site (n)	Prevalence of MG (95% CI)
Durham, NC (n=93)	24.7 (16.0–33.5)
Greensboro, NC (n=152)	38.8 (31.1–46.6)
Pittsburgh, PA (n=174)	27.6 (20.9–34.2)
Birmingham, AL (n=235)	29.8 (23.9–35.6)
New Orleans, LA (n=103)	29.1 (20.4–37.9)
Seattle, WA (n=157)	20.4 (14.1–26.7)
<b>TOTAL (n=914)</b>	<b>28.7 (23.8–33.6)</b>

# M. genitalium screening and diagnostic testing

---

- Population based screening for M. genitalium is not recommended
- Diagnostic testing: NAAT (FDA approved in 2019) for urine, urethral, penile meatal, endocervical, vaginal specimens
- When to test: persistent urethritis that fails initial treatment, also consider for persistent PID or cervicitis


# Over 50-60% of M. genitalium infections have resistance mutations to macrolides (azithro)



National Institutes of Health [HHSN2722013000121,  
HHSN272000010, DIMD16-0039]

Bachmann LH, Kirkcaldy RD, et al. CID 2020

# Resistance guided therapy: *M. genitalium*



# M. genitalium Treatment

Change in 2021 STI Treatment Guidelines

- Sequential treatment for suspected/documentated M. genitalium

Start with Doxycycline to reduce bacterial load

Doxycycline 100 mg  
BID x 7days



Moxifloxacin 400 mg  
BID x 7days

If local macrolide resistance is low or known macrolide sensitive

Doxycycline 100 mg  
BID x 7days




Azithromycin 2.5 gm  
over 4 days

(Azithromycin- 1 gm x 1day then 500 mg x 3day)



# Trichomonas





# **T. Vaginalis screening and diagnostic testing**


---

- Screening for T. vaginalis is recommended for
  - Cis-women with HIV
  - Cis-women in correctional settings
  - Consider for other high prevalence settings
- Diagnostic testing: NAAT for urine, urethral, endocervical (including liquid cytology), vaginal
  - When to test: symptomatic patients

# Treatment Consideration:

## Single dose metronidazole is not as effective as 7 days

- Single dose previously recommended for trich in HIV-negative women, 7-day therapy (500 mg BID) recommended for patients with HIV (CDC TX GL 2015)
- N=623 women randomized 1:1 to single dose MTZ vs 7 day
- Culture TOC, 6-12 days post treatment



Kissinger, 2018 Lancet Infect Dis

# Trichomoniasis Treatment

Change in 2021 STI Treatment  
Guidelines

Recommended regimen: **Vaginal trichomonas (HIV+/HIV-)**

**Metronidazole 500 mg PO BID x 7d**

Metronidazole 2 g PO single dose for men w/ trichomonas or male  
partners)


ACOG 2020 Treatment Guidelines

Metronidazole 500 mg PO BID x 7 d

# Metronidazole and Alcohol

## Change in 2021 STI Treatment Guidelines


- Metronidazole does not actually inhibit acetaldehyde dehydrogenase (as occurs with disulfiram)
- Evidence review: no in vitro or clinical studies, no animal models, and no adverse event reporting
- Refraining from ETOH is unnecessary during treatment



2021 CDC STI Treatment Guidelines  
Fjeld H, Raknes G. Tidsskr Nor Lægeforen. 2014;134(17):1661–3



# Test of Cure vs Retesting



# Test of Cure vs Retesting


TEST OF CURE	Time period	Who
GC (pharynx)	2 weeks	All patients
CT (cervix)	4 weeks	Pregnant patients only
LGV (all sites)	4 weeks	If AZM used instead of DOX (consider)
M. genitalium	3 weeks	If DOX + AZM used instead of MOXI

RETEST FOR REINFECTION	Time period	Who
GC/CT/LGV (all sites)	3 m (anytime from 1-12 m ok)	All patients
Trichomonas	3 m (anytime from 1-12 m ok)	Patients w/vaginal infection





# Syphilis



# Syphilis Diagnosis and Treatment

---

- Neurosyphilis Dx in pts with reactive serology and...
  - Ocular symptoms: if isolated ocular sx, no CN or other neuro involvement, **and** confirmed eye abnormalities on exam, **no CSF exam needed before tx**
  - Ootosyphilis: if isolated auditory abnormalities, CSF likely to be normal, **no CSF needed before tx**
- Follow up: if RPR drops appropriately and patient improves clinically, **no repeat CSF needed for pts without HIV or patients with HIV on ART**
- Treatment: no changes for any stage of syphilis





# Pelvic Inflammatory Disease



# PID Outpatient Treatment: Should Metronidazole be used routinely?

- Randomized Controlled Trial (N=233 cis women)
- Ceftriaxone 250 mg IM plus Doxycycline 100 mg PO BID x 14 days plus
  - Metronidazole 500 mg BID x 14 day OR
  - Placebo BID X 14 day
- Primary outcome: Clinical improvement 3 days
- Additional outcomes: Anerobic organisms in endometrium at 30 days, fever, CMT reduction

# Study Results

---

- Clinical improvement at 3 days similar between two arms
- Metronidazole
  - Reduced anaerobes in endometrium (8% vs 21%,  $p < 0.05$ )
  - Reduced *M. genitalium* (cervical) (4% vs 14%,  $p < 0.05$ )
  - Reduced CMT/pelvic tenderness (9% vs 20%,  $p < 0.05$ )
- **Conclusion: Metronidazole should be routinely added for PID RX**

Wisensfeld et al. CID 2021

# PID IM/Oral Treatment Regimens: Metronidazole for all

Change in 2021 STI  
Treatment Guidelines

## Oral regimens:

- ❖ Ceftriaxone 500 mg IM (or other parenteral 3rd generation cephalosporin) x 1 **or**
- ❖ Cefoxitin 2 g IM **with** probenecid 1 g orally once  
**PLUS**
- ❖ Doxycycline 100 mg orally twice daily for 14 days  
**WITH ~~OR WITHOUT~~**
- ❖ Metronidazole 500 mg orally twice daily for 14 days

# Acknowledgments

---

- Ina Park- UCSF, California Prevention Training Center
- Laura Bachmann- CDC
- Sancta St. Cyr- CDC
- Will Geisler- University of Alabama Birmingham
- Sharon Adler- UCSF, California Prevention Training Center
- Chris Fox- OHSU

Syphilis management? Resistant gonorrhea? STD treatment?

# GOT A TOUGH STD QUESTION?

Get FREE expert STD clinical consultation at your fingertips



 Ask your question

 National STD experts review

 Response within 1-5 business days, depending on urgency

Log on to **[www.STDCCN.org](http://www.STDCCN.org)** for medical professionals nationwide



# Thank you!!



## Any Burning Questions?



Hillary Liss  
[hliss@uw.edu](mailto:hliss@uw.edu)  
206-399-4590

# Acknowledgment

---

The Mountain West AIDS Education and Training (MWAETC) program is supported by the Health Resources and Services Administration (HRSA) of the U.S. Department of Health and Human Services (HHS) as part of an award totaling \$2,990,665 with 0% financed with non-governmental sources.

The content in this presentation are those of the author(s) and do not necessarily represent the official views of, nor an endorsement by, HRSA, HHS, or the U.S. Government.